



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Distant Metastasis in Colorectal Cancer is a Risk Factor for Anastomotic Leakage

Käser, Samuel A ; Mattiello, Diana ; Maurer, Christoph A

DOI: <https://doi.org/10.1245/s10434-015-4941-1>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-135144>

Journal Article

Accepted Version

Originally published at:

Käser, Samuel A; Mattiello, Diana; Maurer, Christoph A (2016). Distant Metastasis in Colorectal Cancer is a Risk Factor for Anastomotic Leakage. *Annals of Surgical Oncology*, 23(3):888-893.

DOI: <https://doi.org/10.1245/s10434-015-4941-1>



Distant metastasis in colorectal cancer is a risk factor for anastomotic leakage

Journal:	<i>Annals of Surgical Oncology</i>
Manuscript ID:	ASO-2015-01-0125.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Käser, Samuel; Hospital of Baselland, Liestal, Department of General, Visceral, Vascular, and Thoracic Surgery Mattiello, Diana; Hospital of Baselland, Liestal, Department of General, Visceral, Vascular, and Thoracic Surgery Maurer, Christoph; Hirslanden Private Clinic Group, Beausite, Surgical Department; Hospital of Baselland, Liestal, Department of General, Visceral, Vascular, and Thoracic Surgery

SCHOLARONE™
Manuscripts

Review

Distant metastasis in colorectal cancer is a risk factor for anastomotic leakage

Käser SA¹, Mattiello D¹, Maurer CA^{1,2}

¹ Department of General, Visceral, Vascular, and Thoracic Surgery, Hospital of Baselland, Rheinstrasse 26, 4410 Liestal, Switzerland

² Hirslanden Private Clinic Group, Beausite, Schänzlihalde 11, 3000 Bern, Switzerland

Correspondence and requests for reprints to:

Christoph A. Maurer, MD, FACS, FRCS (hon.), FEBS

Hirslanden Private Clinic Group, Beausite

Schänzlihalde 11, CH-3000 Bern, Switzerland

Phone: +41 61 906 92 42, Fax: +41 61 906 92 47

E-mail: christoph.maurer@hin.ch

Running head: Stage IV colorectal cancer & anastomotic leaks

The authors declare to have no commercial interests and that there was no financial or material support for this study.

Synopsis: Patients with resection for stage IV colorectal cancer have an increased anastomotic leak rate.

Abstract

Purpose: To investigate if metastatic colorectal cancer (UICC stage IV disease) represents a risk factor for anastomotic leakage after colorectal surgery without major hepatic resection.

Methods: This retrospective cohort study is based on an existing prospective colorectal database with all consecutive colorectal resections done at the authors' institution from 07/2002 to 07/2012 (n=2104). All patients with colorectal resection and primary anastomosis for colorectal cancer were identified (n=500). A temporary loop ileostomy was constructed in low rectal anastomosis up to 6 cm from anal verge (n=128 cases; 26%). Routine contrast enema was done at the occasion of other prospective studies (n=254). UICC stage IV disease was present in n=94 patients (19%), while n=406 (81%) had UICC stage I to III disease.

Results: The overall anastomotic leak rate was 2.6% (13/500), 2.2% (11/500) for both clinical and radiological leaks, and 0.8% (2/254) for radiological leaks only. N=4 were managed conservatively, and n=9 (1.8%) required revision laparotomy. In the case of UICC stage IV disease, the anastomotic leak rate was 6.3% (6/94); in the case of UICC stage I to III disease the leak rate was 1.7% (7/406). UICC stage IV disease (odds ratio 4.4, 95% confidence interval 1.3 - 14.4, p=0.015) and diabetes (odds ratio 5.7, 95% confidence interval 1.7 - 18.7, p=0.004) remained independent risk factors for anastomotic leakage after colorectal surgery.

Conclusions: Patients with stage IV colorectal cancer have an increased anastomotic leak rate after colorectal surgery. Whether this is due to impaired immune system remains speculation.

Introduction

One in five patient with diagnosed colorectal cancer has synchronous liver metastasis and only a minority qualifies for curative liver resection¹. Different treatment strategies in the palliative and in the curative setting are under debate²⁻⁸. However, surprisingly little is known on the specific risks of colorectal surgery in patients with metastatic colorectal cancer.

We aimed to determine if metastatic colorectal cancer (UICC stage IV disease) itself represents a risk factor for anastomotic leakage after colonic or rectal anastomosis without major hepatic resection and to compare it to other known risk factors of anastomotic leakage.

Methods

This retrospective cohort study is based on an existing prospective colorectal database with all consecutive colorectal resections done at the authors' institution from 07/2002 to 07/2012 (n=2104). As different hospitals are located near the authors' institution the study is not population based. Patients with colorectal resections without colonic or rectal anastomosis (n=229) were excluded. From the remaining n=1875 patients, those with the following procedures were excluded: reversal of hartmann's procedure (n=70), redo of anastomosis for anastomotic leak (n=7), colon resection and primary anastomosis for the following diseases: diverticular disease (n=661), colorectal adenoma (n=111), rectal prolapse (n=122), non-colorectal cancer (n=101), complicated appendicitis (n=63), chronic inflammatory bowel disease (n=58), ischemic or infectious colitis (n=44), iatrogenic or traumatic colon perforation (n=37), non-malignant large bowel obstruction (including anastomotic stenosis, radiation injuries, incarcerated hernia n=30), slow-transit constipation (n=22), volvulus (n=11), Ogilvie's syndrome with severe colonic

1
2
3 damage (n=10), colonic bleeding (n=22), and other reasons (n=6). Finally, n=500
4
5 patients with colorectal resection and primary anastomosis for colorectal cancer were
6
7 included in the study.
8

9
10 The routine preoperative workup comprised CT scan of the abdomen and
11
12 conventional X-ray or CT scan of the thorax. In the case of suspicion of liver
13
14 metastasis MRI and intraoperative ultrasound were done.
15

16
17 A temporary loop ileostomy was always constructed in low rectal anastomosis up to 6
18
19 cm from anal verge (n=128 cases; 26%).
20

21
22 The laboratory values measured one day before surgery were assessed.
23

24 Extravasation of water-soluble contrast medium in conventional contrast enema or
25
26 CT scan, abscesses near the colonic or rectal anastomosis without extravasation of
27
28 contrast medium, and anastomotic leaks diagnosed during surgical re-intervention in
29
30 presence of obvious signs of an anastomotic leak such as fecal abdominal drainage
31
32 were accounted as an anastomotic leak. The patients were observed clinically and by
33
34 routine inflammation markers during hospitalization for several days. However, in the
35
36 case of unexpected postoperative course after sending home the patients, they
37
38 reliably were sent back to the present surgical unit for further treatment.
39
40
41

42
43 Routine contrast enema was done at the occasion of other prospective studies
44
45 (n=254), especially in all patients with temporary loop ileostomy (n=131). In the
46
47 remaining patients (n=246) contrast enema was only done in the case of suspicion of
48
49 an anastomotic leak.
50

51
52 UICC stage IV disease was present in n=94 patients (19%), while n=406 (81%) had
53
54 UICC stage I to III disease. An overview of the study methodology is shown in Figure
55
56 1.
57
58
59
60

Statistics: Results are expressed as median and range or mean and standard deviation whenever appropriate. Categorical data was analyzed with the two-sided Fisher's exact test and Pearson's Chi Square if the contingency table was larger than 2x2. Continuous data was analyzed with the Wilcoxon rank sum test. Multivariate analyses were done by logistic regression. P-values < 0.05 were considered statistically significant.

Ethics statement: The present study is a retrospective observational study without any experimental protocols and thus no potential harm for patients.

Results

The only significant difference in the patients' characteristics between patients with stage I-III to patients with stage IV colorectal cancer was the ASA (American Society of Anesthesiologists) score as shown in table 1. The distribution of metastases in patients with metastatic disease (n=94) was: 49% (n=46) liver only, 3% (n=3) lung only, 3% (n=3) peritoneal only, 3% (n=3) retroperitoneal only, 12% (n=11) liver and lung only, 12% (n=11) peritoneal and liver only, 1% (n=1) retroperitoneal and lung only, and 17% (n=16) multiorgan disease.

Patients with metastatic disease were treated with a curative intention in 42.5% (n=40), while 41.5% (n=39) had a planned palliative therapy and the remaining 16% (n=15) had palliative best supportive care due to their poor general state.

The anastomotic leaks were diagnosed on median postoperative day 6 (range 1-15). The overall anastomotic leak rate was 2.6% (13/500). Two of these thirteen anastomotic leaks were diagnosed by routine contrast enema and remained subclinical i.e. without symptoms, two were managed conservatively, and nine required revision laparotomy.

It is noteworthy that three of the six patients with stage IV colorectal cancer who had anastomotic leakage were in a palliative setting.

The analyzed and detected risk factors for the development of an anastomotic leak after resection of colon or rectal cancer (univariate) are shown in Table 2. The results of the multivariate analyses comparing the detected risk factors for the development of an anastomotic leak after resection of colon or rectal cancer are shown in Table 3; stage IV colorectal cancer and diabetes remained statistically significant risk factors for the development of an anastomotic leak, while male gender and resection of left-sided colon cancer or rectal cancer did not.

The leak rate of patients with and without UICC stage IV colorectal cancer is depicted in Figure 2. The leak rate was 9% (6/63) in patients with diabetes mellitus and 1.6% (7/437) without.

Subgroup analyses: colon cancer

From n=339 patients with colon cancer, n=10 had anastomotic leakage (2.9%). With respect to the type of surgery the leak rates were as following: left hemicolectomy 6.7% (8/137); subtotal colectomy (1/31); transverse resection (0/5), right hemicolectomy 0.6% (1/166).

Univariate analysis revealed that stage IV disease ($p=0.006$), left colon cancer ($p=0.010$), and male gender ($p=0.023$) were statistically significant predictors of anastomotic leakage after resection of colon cancer, while the other analyzed factors were not. Multivariate analyses with the detected significant predictors of anastomotic leakage showed that stage IV disease ($p=0.003$, Odds ratio 7.6 [95% confidence interval: 2.0 – 29.4]), left-sided colon cancer ($p=0.025$, Odds ratio 11.1 [95% confidence interval: 1.3 – 91.8]), and male gender ($p=0.045$, Odds ratio 8.7 [95%

confidence interval: 1.0 – 71.5]) remained independent statistically significant predictors of anastomotic leakage after resection of colon cancer.

Subgroup analyses: rectal cancer

From n=161 patients with rectal cancer, n=3 patients had anastomotic leakage (1.9%). In patients with high anterior recto-sigmoid resection with partial mesorectal excision without loop ileostomy for cancer of the upper third of the rectum (n=30) no leak occurred, while in those patients with low anterior rectal resection with loop ileostomy for cancer of the lower two thirds of the rectum (n=128) or with high anterior recto-sigmoid resection with loop ileostomy (n=3) the leak rate was 2.3% (3/131).

Univariate analysis revealed that diabetes (p=0.001) and a lower preoperative hemoglobin value (p=0.023) were statistically significant predictors of anastomotic leakage, while the other analyzed factors were not. Multivariate analyses with the two detected significant predictors of anastomotic leakage showed that diabetes (p<0.001 Odds ratio 7.7E8 [95% confidence interval: 11.9E3 – 5.0E13]) remained a statistically significant predictor of anastomotic leakage while a lower preoperative hemoglobin value (p=0.143 Odds ratio 0.5 [95% confidence interval: 0.2 – 1.3]) did not.

Discussion

Surprisingly little is known on the risks of colorectal surgery in patients with metastatic colorectal cancer. This study shows that metastatic colorectal cancer (UICC stage IV disease) itself is a risk factor for anastomotic leakage after colorectal surgery without major hepatic resection. Is this finding plausible?

1
2
3 First it is known that liver metastasis leads to an altered immune response ^{9,10}.

4
5 Immunosuppression as induced by steroids is a well-known risk factor for the
6
7 development of an anastomotic leak after colorectal surgery ¹¹.

8
9
10 Second up to 65% of the patients with grade IV colorectal cancer have malnutrition
11
12 ¹², and patients with malnutrition also have an altered immune response with an
13
14 exaggerated Interleukin-6 and a suppressed Interleukin-1 response ^{13,14}. A low level
15
16 of Interleukin-1-receptor-antagonist is a known risk factor for adverse outcome after
17
18 colorectal surgery ¹⁵.

19
20 Besides metastatic colorectal cancer as a risk factor for anastomotic leakage,
21
22 diabetes was identified as a second risk factor for anastomotic leakage. Diabetes is a
23
24 known risk factor for delayed wound healing ¹⁸ and thus it can be assumed to be a
25
26 risk factor for anastomotic leakage after colorectal surgery as well. Indeed some
27
28 studies identified diabetes as a risk factor for anastomotic leakage after colorectal
29
30 surgery ^{19,20}, while others did not ^{21,22}. Subgroup analyses revealed that in this study
31
32 diabetes only remained a risk factor for anastomotic leakage in patients with rectal
33
34 cancer. However, further research is needed to confirm if diabetes is a risk factor for
35
36 anastomotic leakage after colorectal surgery or not.

37
38
39
40
41
42
43 Subgroup analyses of those patients with resection of left-sided colon cancer had a
44
45 significant higher leakage rate than those with right-sided colorectal cancer. This is
46
47 known from literature ^{16,17}. We draw the conclusion that the construction of a
48
49 temporary loop ileostomy should be considered in patients with left-sided stage IV
50
51 colon cancer.
52
53
54
55
56
57
58
59
60

Subgroup analyses also identified male gender as a significant risk factor for anastomotic leakage in patients with resection for colon cancer, as known from literature²³.

Limitations

The treatment policy for patients with stage IV colorectal cancer was to do the colorectal resection if feasible and safe. As mentioned above the study is not population based. Thus a patient selection bias with a selection of those patients in a better general state is possible. However, if more patients in a worse general state with an expected higher leakage rate would have been included, it can be assumed that stage IV disease would have turned out to be an even stronger predictor of anastomotic leakage.

As only about half of the patients were investigated by routine radiology (n=254) few subclinical anastomotic leaks could have been missed in those patients who did not have routine radiology (n=246). As one of the two patients with a subclinical leak had stage I cancer and the other had stage IV cancer a bias is not expected.

This study has a retrospective design. Thus, a further bias cannot be excluded. However, a bias was minimized by including all consecutive patients registered in a prospective database who had colorectal resection at the same institution in a predefined period of ten years.

To confirm the results of this study and to prove the value of a routine temporary loop ileostomy in patients with left-sided stage IV colon cancer a prospective randomized trial is required.

Conclusion

Patients with stage IV colorectal cancer have an increased leak rate after colonic or rectal anastomosis. Whether this is due to impaired immune system remains speculation.

For Peer Review

Table 1 Patients’ characteristics

Risk factor	Stage I - III colorectal cancer (n=406)	Stage IV colorectal cancer (n=94)	p-value
Mean age (SD)	70 (12) years	69 (12) years	0.190
Male gender	n=237/406	n=53/94	0.729
Median ASA Score (Range)	2 (1-5)	3 (1-4)	0.012
Cardiac comorbidities	n=204/356 (57%)	n=55/85 (65%)	0.372
Vascular comorbidities	n=41/342 (12%)	n=6/85 (7%)	0.327
Diabetes	n=54/356 (15%)	n=9/85 (11%)	0.388
Pulmonary comorbidities	n=55/356 (15%)	n=18/85 (21%)	0.198
Tobacco abuse	n=48/356 (13%)	n=15/85 (18%)	0.307
Obesity	n=195/356 (55%)	n=54/85 (64%)	0.180
Dementia	n=12/355 (3%)	n=0/85 (0%)	0.135
Immunosuppressive drugs	n=6/352 (2%)	n=2/85 (2%)	0.656
Non-steroidal anti- inflammatory drugs	n=62/352 (18%)	n=13/85 (15%)	0.749
Presence of peritonitis	n=22/329 (7%)	n=5/85 (6%)	1.000
Large bowel obstruction	n=37/351 (11%)	n=10/85 (12%)	0.700
Urgent surgery	n=49/351 (14%)	n=12/85 (15%)	1.000

SD: standard deviation, ASA: American Society of Anesthesiologists

Table 2 Results of the univariate analyses of the risk factors of anastomotic leakage after colorectal surgery for colorectal cancer.

Risk factor	Anastomotic leak (n=13)	No anastomotic leak (n=487)	p-value
Mean age (SD)	73 (8) years	70 (12) years	0.516
Male gender	12/13	278/487 (57%)	0.010
Median ASA Score (Range)	2 (1-3)	2 (1-5)	0.657
Cardiac comorbidities	9/13	250/428 (58%)	0.584
Vascular comorbidities	2/13	45/428 (11%)	0.638
Diabetes	6/13	57/428 (13%)	0.005
Pulmonary comorbidities	1/13	72/428 (17%)	0.704
Tobacco abuse	2/13	61/428 (14%)	1.000
Obesity	8/13	241/428 (56%)	0.783
Dementia	0/13	12/427 (3%)	1.000
Immunosuppressive drugs	0/13	8/424 (2%)	1.000
Non-steroidal anti- inflammatory drugs	3/13	72/424 (17%)	0.475
Preoperative radio- and/or chemotherapy	0/13	43/472 (9%)	0.617
Presence of peritonitis	0/13	27/396 (7%)	1.000
Large bowel obstruction	1/13	46/377 (12%)	1.000
Urgent surgery	1/13	60/396 (15%)	0.702
ypT0 or T1 cancer	2/13	52/487 (11%)	

T2 cancer	4/13	73/487 (15%)	
T3 cancer	3/13	245/487 (50%)	
T4 cancer	4/13	117/487 (24%)	0.218
Nodal positive cancer	8/13	219/473 (46%)	0.270
Stage IV colorectal cancer	6/13	88/487 (18%)	0.021
Resection of left-sided colon cancer or rectal cancer	12/13	317/487 (65%)	0.0417
Preoperative mean (SD) white blood cell count (1/10E9)	8.0 (3.5)	7.7 (3.0)	0.629
Preoperative mean (SD) Hemoglobin	12.6 (2.0)	12.8 (2.1)	0.814
Preoperative mean (SD) Sodium	138.5 (2.9)	139.7 (3.1)	0.0935
Preoperative mean (SD) Potassium	4.3 (0.4)	4.1 (0.4)	0.324
Preoperative mean (SD) Creatinine	82.8 (23.8)	79.7 (40.8)	0.328
Preoperative mean (SD) Albumin	38 (4.0)	39.3 (5.5)	0.1872

SD: standard deviation, ASA: American Society of Anesthesiologists

Table 3 Results of the multivariate analyses of the detected risk factors for anastomotic leakage after colorectal surgery for colorectal cancer.

Risk factor	Odds ratio	95%confidence interval	p-value
Diabetes	5.7	1.7 – 18.7	0.004
Stage IV colorectal cancer	4.4	1.3 – 14.4	0.015
Male gender	7.8	1.0 – 62.6	0.052
Resection of left-sided colorectal cancer or rectal cancer	6.7	0.8 – 53.0	0.073

ASA: American Society of Anesthesiologists

Figure 1: Flow diagram of the study methodology

Figure 2: Anastomotic leakage rate in Stage I to III versus Stage IV colorectal cancer

References

1. van der Pool AE, Damhuis RA, Ijzermans JN, et al. Trends in incidence, treatment and survival of patients with stage IV colorectal cancer: a population-based series. *Colorectal Dis* 2012; 14:56-61.

2. Chen GQ, Li J, Ding KF. A meta-analysis of the safety of simultaneous versus staged resection for synchronous liver metastasis from colorectal cancer. *Zhonghua wei chang wai ke za zhi* 2010; 13:337-341.

3. Hillingso JG, Wille-Jorgensen P. Staged or simultaneous resection of synchronous liver metastases from colorectal cancer--a systematic review. *Colorectal Dis* 2009; 11:3-10.

4. Hopt UT, Drogitz O, Neeff H. Timing of resection in patients with colorectal carcinoma and synchronous liver metastases. *Zentralbl Chir* 2009; 134:425-429.

5. de Mestier L, Manceau G, Neuzillet C, et al. Primary tumor resection in colorectal cancer with unresectable synchronous metastases: A review. *World J Gastrointest Oncol* 2014; 6:156-169.


6. Rosen SA, Buell JF, Yoshida A, et al. Initial presentation with stage IV colorectal cancer: how aggressive should we be? *Arch Surg* 2000; 135:530-534.

7. Vibert E, Bretagnol F, Alves A, Pocard M, Valleur P, Panis Y. Multivariate analysis of predictive factors for early postoperative death after colorectal surgery in patients with colorectal cancer and synchronous unresectable liver metastases. *Dis Colon Rectum* 2007; 50:1776-1782.


8. Ruo L, Gougoutas C, Paty PB, Guillem JG, Cohen AM, Wong WD. Elective bowel resection for incurable stage IV colorectal cancer: prognostic variables for asymptomatic patients. *J Am Coll Surg* 2003; 196:722-728.

- 1
2
3 9. Brudvik KW, Henjum K, Aandahl EM, Bjornbeth BA, Tasken K. Regulatory T-cell-
4 mediated inhibition of antitumor immune responses is associated with clinical
5 outcome in patients with liver metastasis from colorectal cancer. *Cancer*
6 *Immunol Immunother* 2012; 61:1045-1053.
- 7
8
9
10
11 10. Giakoustidis A, Mudan S, Hagemann T. Tumour Microenvironment: Overview
12 with an Emphasis on the Colorectal Liver Metastasis Pathway. *Cancer*
13 *Microenviron* 2014 [Epub ahead of print].
- 14
15
16
17
18 11. Konishi T, Watanabe T, Kishimoto J, Nagawa H. Risk factors for anastomotic
19 leakage after surgery for colorectal cancer: results of prospective surveillance. *J*
20 *Am Coll Surg* 2006; 202:439-444.
- 21
22
23
24
25 12. Barret M, Malka D, Aparicio T, et al. Nutritional status affects treatment
26 tolerability and survival in metastatic colorectal cancer patients: results of an
27 AGEO prospective multicenter study. *Oncology* 2011; 81:395-402.
- 28
29
30
31
32 13. Hatada T, Miki C. Nutritional status and postoperative cytokine response in
33 colorectal cancer patients. *Cytokine* 2000; 12:1331-1336.
- 34
35
36
37 14. Nakagoe T, Tsuji T, Sawai T, et al. Increased serum levels of interleukin-6 in
38 malnourished patients with colorectal cancer. *Cancer Lett* 2003; 202:109-115.
- 39
40
41
42 15. Miki C, Inoue Y, Toiyama Y, et al. Deficiency in systemic interleukin-1 receptor
43 antagonist production as an operative risk factor in malnourished elderly
44 patients with colorectal carcinoma. *Crit Care Med* 2005; 33:177-180.
- 45
46
47
48 16. Bakker IS, Grossmann I, Henneman D, Havenga K, Wiggers T. Risk factors for
49 anastomotic leakage and leak-related mortality after colonic cancer surgery in a
50 nationwide audit. *Br J Surg* 2014; 101:424-432
- 51
52
53
54
55
56
57
58
59
60

1
2
3 17. Krarup PM, Jorgensen LN, Andreassen AH, Harling H, Danish Colorectal Cancer G.
4 A nationwide study on anastomotic leakage after colonic cancer surgery.
5 Colorectal Dis 2012; 14: 661-667.
6
7
8
9
10 18. Eming SA, Martin P, Tomic-Canic M. Wound repair and regeneration:
11 Mechanisms, signaling, and translation. Sci Transl Med 2014; 6:265-266.
12
13
14 19. Cong ZJ, Fu CG, Wang HT, Liu LJ, Zhang W, Wang H. Influencing factors of
15 symptomatic anastomotic leakage after anterior resection of the rectum for
16 cancer. World J Surg 2009; 33:1292-1297.
17
18
19
20
21 20. Volk A, Kersting S, Held HC, Saeger HD. Risk factors for morbidity and mortality
22 after single-layer continuous suture for ileocolonic anastomosis. Int J Colorectal
23 Dis 2011; 26:321-327.
24
25
26
27
28 21. Liu Y, Wan X, Wang G, et al. A scoring system to predict the risk of anastomotic
29 leakage after anterior resection for rectal cancer. J Surg Oncol 2014; 109:122-
30 125.
31
32
33
34 22. Luna-Perez P, Rodriguez-Ramirez SE, Gutierrez de la Barrera M, Labastida S.
35 Multivariate analysis of risk factors associated with dehiscence of colorectal
36 anastomosis after anterior or lower anterior resection for sigmoid or rectal
37 cancer. Rev Invest Clin 2002; 54:501-508.
38
39
40
41
42
43 23. Lipska MA, Bissett IP, Parry BR, Merrie AE. Anastomotic leakage after lower
44 gastrointestinal anastomosis: men are at a higher risk. ANZ J Surg 2006; 76:579-
45 585.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Flow diagram of the study methodology
435x284mm (72 x 72 DPI)



Anastomotic leakage rate in Stage I to III versus Stage IV colorectal cancer
360x299mm (120 x 120 DPI)

Distant metastasis in colorectal cancer is a risk factor for anastomotic leakage

Käser SA¹, Mattiello D¹, Maurer CA^{1,2}

¹ Department of General, Visceral, Vascular, and Thoracic Surgery, Hospital of Baselland, Rheinstrasse 26, 4410 Liestal, Switzerland

² Hirslanden Private Clinic Group, Beausite, Schänzlihalde 11, 3000 Bern, Switzerland

Correspondence and requests for reprints to:

Christoph A. Maurer, MD, FACS, FRCS (hon.), FEBS

Hirslanden Private Clinic Group, Beausite

Schänzlihalde 11, CH-3000 Bern, Switzerland

Phone: +41 61 906 92 42, Fax: +41 61 906 92 47

E-mail: christoph.maurer@hin.ch

Running head: Stage IV colorectal cancer & anastomotic leaks

The authors declare to have no commercial interests and that there was no financial or material support for this study.

Synopsis: Patients with resection for stage IV colorectal cancer have an increased anastomotic leak rate.

Abstract

Purpose: To investigate if metastatic colorectal cancer (UICC stage IV disease) represents a risk factor for anastomotic leakage after colorectal surgery without major hepatic resection.

Methods: This retrospective cohort study is based on an existing prospective colorectal database with all consecutive colorectal resections done at the authors' institution from 07/2002 to 07/2012 (n=2104). All patients with colorectal resection and primary anastomosis for colorectal cancer were identified (n=500). A temporary loop ileostomy was constructed in low rectal anastomosis up to 6 cm from anal verge (n=128 cases; 26%). Routine contrast enema was done at the occasion of other prospective studies (n=254). UICC stage IV disease was present in n=94 patients (19%), while n=406 (81%) had UICC stage I to III disease.

Results: **The overall anastomotic leak rate was 2.6% (13/500), 2.2% (11/500) for both clinical and radiological leaks, and 0.8% (2/254) for radiological leaks only.** N=4 were managed conservatively, and n=9 (1.8%) required revision laparotomy. In the case of UICC stage IV disease, the anastomotic leak rate was 6.3% (6/94); in the case of UICC stage I to III disease the leak rate was 1.7% (7/406). **UICC stage IV disease (odds ratio 4.4, 95% confidence interval 1.3 - 14.4, p=0.015) and diabetes (odds ratio 5.7, 95% confidence interval 1.7 - 18.7, p=0.004) remained independent risk factors for anastomotic leakage after colorectal surgery.**

Conclusions: Patients with stage IV colorectal cancer have an increased anastomotic leak rate after colorectal surgery. Whether this is due to impaired immune system remains speculation.

For Peer Review

Introduction

One in five patient with diagnosed colorectal cancer has synchronous liver metastasis and only a minority qualifies for curative liver resection¹. Different treatment strategies in the palliative and in the curative setting are under debate²⁻⁸. However, surprisingly little is known on the specific risks of colorectal surgery in patients with metastatic colorectal cancer. We aimed to determine if metastatic colorectal cancer (UICC stage IV disease) itself represents a risk factor for anastomotic leakage after colonic or rectal anastomosis without major hepatic resection and to compare it to other known risk factors of anastomotic leakage.

Methods

This retrospective cohort study is based on an existing prospective colorectal database with all consecutive colorectal resections done at the authors' institution from 07/2002 to 07/2012 (n=2104). As different hospitals are located near the authors' institution the study is not population based. Patients with colorectal resections without colonic or rectal anastomosis (n=229) were excluded. From the remaining n=1875 patients, those with the following procedures were excluded: reversal of hartmann's procedure (n=70), redo of anastomosis for anastomotic leak (n=7), colon resection and primary anastomosis for the following diseases: diverticular disease (n=661), colorectal adenoma (n=111), rectal prolapse (n=122), non-colorectal cancer (n=101), complicated appendicitis (n=63), chronic inflammatory bowel disease (n=58), ischemic or infectious colitis (n=44), iatrogenic or traumatic colon perforation (n=37), non-malignant large bowel obstruction (including anastomotic stenosis, radiation injuries, incarcerated hernia n=30), slow-transit

constipation (n=22), volvulus (n=11), Ogilvie's syndrome with severe colonic damage (n=10), colonic bleeding (n=22), and other reasons (n=6). Finally, n=500 patients with colorectal resection and primary anastomosis for colorectal cancer were included in the study.

The routine preoperative workup comprised CT scan of the abdomen and conventional X-ray or CT scan of the thorax. In the case of suspicion of liver metastasis MRI and intraoperative ultrasound were done.

A temporary loop ileostomy was always constructed in low rectal anastomosis up to 6 cm from anal verge (n=128 cases; 26%).

The laboratory values measured one day before surgery were assessed.

Extravasation of water-soluble contrast medium in conventional contrast enema or CT scan, abscesses near the colonic or rectal anastomosis without extravasation of contrast medium, and anastomotic leaks diagnosed during surgical re-intervention in presence of obvious signs of an anastomotic leak such as fecal abdominal drainage were accounted as an anastomotic leak. The patients were observed clinically and by routine inflammation markers during hospitalization for several days. However, in the case of unexpected postoperative course after sending home the patients, they reliably were sent back to the present surgical unit for further treatment.

Routine contrast enema was done at the occasion of other prospective studies (n=254), especially in all patients with temporary loop ileostomy (n=131). In the remaining patients (n=246) contrast enema was only done in the case of suspicion of an anastomotic leak.

UICC stage IV disease was present in n=94 patients (19%), while n=406 (81%) had UICC stage I to III disease. An overview of the study methodology is shown in Figure 1.

Statistics: Results are expressed as median and range or mean and standard deviation whenever appropriate. Categorical data was analyzed with the two-sided Fisher's exact test and Pearson's Chi Square if the contingency table was larger than 2x2. Continuous data was analyzed with the Wilcoxon rank sum test. Multivariate analyses were done by logistic regression. P-values < 0.05 were considered statistically significant.

Ethics statement: The present study is a retrospective observational study without any experimental protocols and thus no potential harm for patients.

Results

The only significant difference in the patients' characteristics between patients with stage I-III to patients with stage IV colorectal cancer was the ASA (American Society of Anesthesiologists) score as shown in table 1. The distribution of metastases in patients with metastatic disease (n=94) was: 49% (n=46) liver only, 3% (n=3) lung only, 3% (n=3) peritoneal only, 3% (n=3) retroperitoneal only, 12% (n=11) liver and lung only, 12% (n=11) peritoneal and liver only, 1% (n=1) retroperitoneal and lung only, and 17% (n=16) multiorgan disease.

Patients with metastatic disease were treated with a curative intention in 42.5% (n=40), while 41.5% (n=39) had a planned palliative therapy and the remaining 16% (n=15) had palliative best supportive care due to their poor general state.

The anastomotic leaks were diagnosed on median postoperative day 6 (range 1-15). The overall anastomotic leak rate was 2.6% (13/500). Two of these thirteen anastomotic leaks were diagnosed by routine contrast enema and remained subclinical i.e. without symptoms, two were managed conservatively, and nine required revision laparotomy.

It is noteworthy that three of the six patients with stage IV colorectal cancer who had anastomotic leakage were in a palliative setting.

The analyzed and detected risk factors for the development of an anastomotic leak after resection of colon or rectal cancer (univariate) are shown in Table 2. The results of the multivariate analyses comparing the detected risk factors for the development of an anastomotic leak after resection of colon or rectal cancer are shown in Table 3; stage IV colorectal cancer and diabetes remained statistically significant risk factors for the development of an anastomotic leak, while male gender and resection of left-sided colon cancer or rectal cancer did not.

The leak rate of patients with and without UICC stage IV colorectal cancer is depicted in Figure 2. The leak rate was 9% (6/63) in patients with diabetes mellitus and 1.6% (7/437) without.

Subgroup analyses: colon cancer

From n=339 patients with colon cancer, n=10 had anastomotic leakage (2.9%). With respect to the type of surgery the leak rates were as following: left hemicolectomy 6.7% (8/137); subtotal colectomy (1/31); transverse resection (0/5), right hemicolectomy 0.6% (1/166).

Univariate analysis revealed that stage IV disease ($p=0.006$), left colon cancer ($p=0.010$), and male gender ($p=0.023$) were statistically significant predictors of anastomotic leakage after resection of colon cancer, while the other analyzed factors were not. Multivariate analyses with the detected significant predictors of anastomotic leakage showed that stage IV disease ($p=0.003$, Odds ratio 7.6 [95% confidence interval: 2.0 – 29.4]), left-sided colon cancer ($p=0.025$, Odds ratio 11.1 [95% confidence interval: 1.3 – 91.8]), and male gender ($p=0.045$, Odds ratio 8.7 [95%

confidence interval: 1.0 – 71.5]) remained independent statistically significant predictors of anastomotic leakage after resection of colon cancer.

Subgroup analyses: rectal cancer

From n=161 patients with rectal cancer, n=3 patients had anastomotic leakage (1.9%). In patients with high anterior recto-sigmoid resection with partial mesorectal excision without loop ileostomy for cancer of the upper third of the rectum (n=30) no leak occurred, while in those patients with low anterior rectal resection with loop ileostomy for cancer of the lower two thirds of the rectum (n=128) or with high anterior recto-sigmoid resection with loop ileostomy (n=3) the leak rate was 2.3% (3/131).

Univariate analysis revealed that diabetes (p=0.001) and a lower preoperative hemoglobin value (p=0.023) were statistically significant predictors of anastomotic leakage, while the other analyzed factors were not. Multivariate analyses with the two detected significant predictors of anastomotic leakage showed that diabetes (p<0.001 Odds ratio 7.7E8 [95% confidence interval: 11.9E3 – 5.0E13]) remained a statistically significant predictor of anastomotic leakage while a lower preoperative hemoglobin value (p=0.143 Odds ratio 0.5 [95% confidence interval: 0.2 – 1.3]) did not.

Discussion

Surprisingly little is known on the risks of colorectal surgery in patients with metastatic colorectal cancer. This study shows that metastatic colorectal cancer (UICC stage IV disease) itself is a risk factor for anastomotic leakage after colorectal surgery without major hepatic resection. Is this finding plausible?

First it is known that liver metastasis leads to an altered immune response^{9,10}.

Immunosuppression as induced by steroids is a well-known risk factor for the development of an anastomotic leak after colorectal surgery¹¹.

Second up to 65% of the patients with grade IV colorectal cancer have malnutrition¹², and patients with malnutrition also have an altered immune response with an exaggerated Interleukin-6 and a suppressed Interleukin-1 response^{13,14}. A low level of Interleukin-1-receptor-antagonist is a known risk factor for adverse outcome after colorectal surgery¹⁵.

Besides metastatic colorectal cancer as a risk factor for anastomotic leakage, diabetes was identified as a second risk factor for anastomotic leakage. Diabetes is a known risk factor for delayed wound healing¹⁸ and thus it can be assumed to be a risk factor for anastomotic leakage after colorectal surgery as well. Indeed some studies identified diabetes as a risk factor for anastomotic leakage after colorectal surgery^{19,20}, while others did not^{21,22}. Subgroup analyses revealed that in this study diabetes only remained a risk factor for anastomotic leakage in patients with rectal cancer. However, further research is needed to confirm if diabetes is a risk factor for anastomotic leakage after colorectal surgery or not.

Subgroup analyses of those patients with resection of left-sided colon cancer had a significant higher leakage rate than those with right-sided colorectal cancer. This is known from literature^{16,17}. We draw the conclusion that the construction of a temporary loop ileostomy should be considered in patients with left-sided stage IV colon cancer.

Subgroup analyses also identified male gender as a significant risk factor for anastomotic leakage in patients with resection for colon cancer, as known from literature²³.

Limitations

The treatment policy for patients with stage IV colorectal cancer was to do the colorectal resection if feasible and safe. As mentioned above the study is not population based. Thus a patient selection bias with a selection of those patients in a better general state is possible. However, if more patients in a worse general state with an expected higher leakage rate would have been included, it can be assumed that stage IV disease would have turned out to be an even stronger predictor of anastomotic leakage.

As only about half of the patients were investigated by routine radiology (n=254) few subclinical anastomotic leaks could have been missed in those patients who did not have routine radiology (n=246). As one of the two patients with a subclinical leak had stage I cancer and the other had stage IV cancer a bias is not expected.

This study has a retrospective design. Thus, a further bias cannot be excluded. However, a bias was minimized by including all consecutive patients registered in a prospective database who had colorectal resection at the same institution in a predefined period of ten years.

To confirm the results of this study and to prove the value of a routine temporary loop ileostomy in patients with left-sided stage IV colon cancer a prospective randomized trial is required.

Conclusion

Patients with stage IV colorectal cancer have an increased leak rate after colonic or rectal anastomosis. Whether this is due to impaired immune system remains speculation.

For Peer Review

Table 1 Patients’ characteristics

Risk factor	Stage I - III colorectal cancer (n=406)	Stage IV colorectal cancer (n=94)	p-value
Mean age (SD)	70 (12) years	69 (12) years	0.190
Male gender	n=237/406	n=53/94	0.729
Median ASA Score (Range)	2 (1-5)	3 (1-4)	0.012
Cardiac comorbidities	n=204/356 (57%)	n=55/85 (65%)	0.372
Vascular comorbidities	n=41/342 (12%)	n=6/85 (7%)	0.327
Diabetes	n=54/356 (15%)	n=9/85 (11%)	0.388
Pulmonary comorbidities	n=55/356 (15%)	n=18/85 (21%)	0.198
Tobacco abuse	n=48/356 (13%)	n=15/85 (18%)	0.307
Obesity	n=195/356 (55%)	n=54/85 (64%)	0.180
Dementia	n=12/355 (3%)	n=0/85 (0%)	0.135
Immunosuppressive drugs	n=6/352 (2%)	n=2/85 (2%)	0.656
Non-steroidal anti- inflammatory drugs	n=62/352 (18%)	n=13/85 (15%)	0.749
Presence of peritonitis	n=22/329 (7%)	n=5/85 (6%)	1.000
Large bowel obstruction	n=37/351 (11%)	n=10/85 (12%)	0.700
Urgent surgery	n=49/351 (14%)	n=12/85 (15%)	1.000

SD: standard deviation, ASA: American Society of Anesthesiologists

Table 2 Results of the univariate analyses of the risk factors of anastomotic leakage after colorectal surgery for colorectal cancer.

Risk factor	Anastomotic leak (n=13)	No anastomotic leak (n=487)	p-value
Mean age (SD)	73 (8) years	70 (12) years	0.516
Male gender	12/13	278/487 (57%)	0.010
Median ASA Score (Range)	2 (1-3)	2 (1-5)	0.657
Cardiac comorbidities	9/13	250/428 (58%)	0.584
Vascular comorbidities	2/13	45/428 (11%)	0.638
Diabetes	6/13	57/428 (13%)	0.005
Pulmonary comorbidities	1/13	72/428 (17%)	0.704
Tobacco abuse	2/13	61/428 (14%)	1.000
Obesity	8/13	241/428 (56%)	0.783
Dementia	0/13	12/427 (3%)	1.000
Immunosuppressive drugs	0/13	8/424 (2%)	1.000
Non-steroidal anti- inflammatory drugs	3/13	72/424 (17%)	0.475
Preoperative radio- and/or chemotherapy	0/13	43/472 (9%)	0.617
Presence of peritonitis	0/13	27/396 (7%)	1.000
Large bowel obstruction	1/13	46/377 (12%)	1.000
Urgent surgery	1/13	60/396 (15%)	0.702
ypT0 or T1 cancer	2/13	52/487 (11%)	

T2 cancer	4/13	73/487 (15%)	
T3 cancer	3/13	245/487 (50%)	
T4 cancer	4/13	117/487 (24%)	0.218
Nodal positive cancer	8/13	219/473 (46%)	0.270
Stage IV colorectal cancer	6/13	88/487 (18%)	0.021
Resection of left-sided colon cancer or rectal cancer	12/13	317/487 (65%)	0.0417
Preoperative mean (SD) white blood cell count (1/10E9)	8.0 (3.5)	7.7 (3.0)	0.629
Preoperative mean (SD) Hemoglobin	12.6 (2.0)	12.8 (2.1)	0.814
Preoperative mean (SD) Sodium	138.5 (2.9)	139.7 (3.1)	0.0935
Preoperative mean (SD) Potassium	4.3 (0.4)	4.1 (0.4)	0.324
Preoperative mean (SD) Creatinine	82.8 (23.8)	79.7 (40.8)	0.328
Preoperative mean (SD) Albumin	38 (4.0)	39.3 (5.5)	0.1872

SD: standard deviation, ASA: American Society of Anesthesiologists

Table 3 Results of the multivariate analyses of the detected risk factors for anastomotic leakage after colorectal surgery for colorectal cancer.

Risk factor	Odds ratio	95%confidence interval	p-value
Diabetes	5.7	1.7 – 18.7	0.004
Stage IV colorectal cancer	4.4	1.3 – 14.4	0.015
Male gender	7.8	1.0 – 62.6	0.052
Resection of left-sided colorectal cancer or rectal cancer	6.7	0.8 – 53.0	0.073

ASA: American Society of Anesthesiologists

Figure 1: Flow diagram of the study methodology

Figure 2: Anastomotic leakage rate in Stage I to III versus Stage IV colorectal cancer

References

1. van der Pool AE, Damhuis RA, Ijzermans JN, et al. Trends in incidence, treatment and survival of patients with stage IV colorectal cancer: a population-based series. *Colorectal Dis* 2012; 14:56-61.

2. Chen GQ, Li J, Ding KF. A meta-analysis of the safety of simultaneous versus staged resection for synchronous liver metastasis from colorectal cancer. *Zhonghua wei chang wai ke za zhi* 2010; 13:337-341.

3. Hillingso JG, Wille-Jorgensen P. Staged or simultaneous resection of synchronous liver metastases from colorectal cancer--a systematic review. *Colorectal Dis* 2009; 11:3-10.

4. Hopt UT, Drogitz O, Neeff H. Timing of resection in patients with colorectal carcinoma and synchronous liver metastases. *Zentralbl Chir* 2009; 134:425-429.

5. de Mestier L, Manceau G, Neuzillet C, et al. Primary tumor resection in colorectal cancer with unresectable synchronous metastases: A review. *World J Gastrointest Oncol* 2014; 6:156-169.

6. Rosen SA, Buell JF, Yoshida A, et al. Initial presentation with stage IV colorectal cancer: how aggressive should we be? *Arch Surg* 2000; 135:530-534.

7. Vibert E, Bretagnol F, Alves A, Pocard M, Valleur P, Panis Y. Multivariate analysis of predictive factors for early postoperative death after colorectal surgery in patients with colorectal cancer and synchronous unresectable liver metastases. *Dis Colon Rectum* 2007; 50:1776-1782.

8. Ruo L, Gougoutas C, Paty PB, Guillem JG, Cohen AM, Wong WD. Elective bowel resection for incurable stage IV colorectal cancer: prognostic variables for asymptomatic patients. *J Am Coll Surg* 2003; 196:722-728.

- 1
2
3 9. Brudvik KW, Henjum K, Aandahl EM, Bjornbeth BA, Tasken K. Regulatory T-cell-
4 mediated inhibition of antitumor immune responses is associated with clinical
5 outcome in patients with liver metastasis from colorectal cancer. *Cancer*
6 *Immunol Immunother* 2012; 61:1045-1053.
7
8
9
10
- 11 10. Giakoustidis A, Mudan S, Hagemann T. Tumour Microenvironment: Overview
12 with an Emphasis on the Colorectal Liver Metastasis Pathway. *Cancer*
13 *Microenviron* 2014 [Epub ahead of print].
14
15
16
17
- 18 11. Konishi T, Watanabe T, Kishimoto J, Nagawa H. Risk factors for anastomotic
19 leakage after surgery for colorectal cancer: results of prospective surveillance. *J*
20 *Am Coll Surg* 2006; 202:439-444.
21
22
23
24
- 25 12. Barret M, Malka D, Aparicio T, et al. Nutritional status affects treatment
26 tolerability and survival in metastatic colorectal cancer patients: results of an
27 AGEO prospective multicenter study. *Oncology* 2011; 81:395-402.
28
29
30
31
- 32 13. Hatada T, Miki C. Nutritional status and postoperative cytokine response in
33 colorectal cancer patients. *Cytokine* 2000; 12:1331-1336.
34
35
36
- 37 14. Nakagoe T, Tsuji T, Sawai T, et al. Increased serum levels of interleukin-6 in
38 malnourished patients with colorectal cancer. *Cancer Lett* 2003; 202:109-115.
39
40
41
- 42 15. Miki C, Inoue Y, Toiyama Y, et al. Deficiency in systemic interleukin-1 receptor
43 antagonist production as an operative risk factor in malnourished elderly
44 patients with colorectal carcinoma. *Crit Care Med* 2005; 33:177-180.
45
46
47
- 48 16. Bakker IS, Grossmann I, Henneman D, Havenga K, Wiggers T. Risk factors for
49 anastomotic leakage and leak-related mortality after colonic cancer surgery in a
50 nationwide audit. *Br J Surg* 2014; 101:424-432
51
52
53
54
55
56
57
58
59
60

1
2
3 17. Krarup PM, Jorgensen LN, Andreasen AH, Harling H, Danish Colorectal Cancer G.
4 A nationwide study on anastomotic leakage after colonic cancer surgery.
5 Colorectal Dis 2012; 14: 661-667.
6
7
8
9
10 18. Eming SA, Martin P, Tomic-Canic M. Wound repair and regeneration:
11 Mechanisms, signaling, and translation. Sci Transl Med 2014; 6:265-266.
12
13
14 19. Cong ZJ, Fu CG, Wang HT, Liu LJ, Zhang W, Wang H. Influencing factors of
15 symptomatic anastomotic leakage after anterior resection of the rectum for
16 cancer. World J Surg 2009; 33:1292-1297.
17
18
19
20
21 20. Volk A, Kersting S, Held HC, Saeger HD. Risk factors for morbidity and mortality
22 after single-layer continuous suture for ileocolonic anastomosis. Int J Colorectal
23 Dis 2011; 26:321-327.
24
25
26
27
28 21. Liu Y, Wan X, Wang G, et al. A scoring system to predict the risk of anastomotic
29 leakage after anterior resection for rectal cancer. J Surg Oncol 2014; 109:122-
30 125.
31
32
33
34 22. Luna-Perez P, Rodriguez-Ramirez SE, Gutierrez de la Barrera M, Labastida S.
35 Multivariate analysis of risk factors associated with dehiscence of colorectal
36 anastomosis after anterior or lower anterior resection for sigmoid or rectal
37 cancer. Rev Invest Clin 2002; 54:501-508.
38
39
40
41
42
43 23. Lipska MA, Bissett IP, Parry BR, Merrie AE. Anastomotic leakage after lower
44 gastrointestinal anastomosis: men are at a higher risk. ANZ J Surg 2006; 76:579-
45 585.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60